AD-A252 021





Quarterly Report No. 5 • June 1992

DEVELOPMENT OF A THERAPEUTIC AGENT FOR WOUND-HEALING ENHANCEMENT

Amrit K. Judd, Ph. D., Senior Peptide Chemist Bio-Organic Chemistry Laboratory

SRI Project LSU-2323

Prepared for:

Naval Medical R&D Command National Naval Medical Center Combat Casualty Care, Research Area Manager Building 1 (Tower Bldg.), 11th Floor 8901 Wisconsin Avenue Bethesda, Maryland 20889-5044

Attn: Capt. Stephen B. Lewis, Scientific Officer

Contract No. N00014-91-C-0066

Approved:

Masato Tanabe, Ph.D. Director Bio-Organic Chemistry Laboratory This document has been approved for public release and sale; its distribution is unlimited.

92-16231

92 6 18 018

SCIENTIFIC PROGRESS DURING THIS QUARTER

Peptide Synthesis

During this quarter we prepared four peptides, as listed below:

PDGF A (53-63)	Peptide #5, Table 3, p. 18 in the original proposal
PDGF A (66-78)	Peptide #6, Table 3, p. 18 in the original proposal
PDGF A (77-88)	Peptide #7, Table 3, p. 18 in the original proposal
PDGF B (25-42)	Peptide #2, Table 4, p. 18 in the original proposal

Biological Assays

During this quarter we conducted cell-binding experiments and assays for mitogenesis and chemotaxis on the four peptides mentioned above. The results are shown in Tables 1 through 3. None of these peptides showed any cell binding or mitogenesis. PDGF A (77-88) showed chemotaxis. We are repeating the assay for this peptide.

Table 1						
RESULTS	OF	COMPETITIVE	BINDING	ASSAY		

Peptide	Concentration (ng or µg per weil)	СРМ	Average ± SD	Maximum Binding	Inhibition (%)
125 _I PDGF	5 ng	37,890 33,988 38,143	36,674 ± 2,329	100	0
PDGF BB Accesion For	50 ng	19,689 21,135 18,053	19,626 ± 1,542	54	46
NTIS CRA&I DTIC TAB Unannounced	25 ng	22,427 22,049 23,350	22,609 ± 669	62	38
By pen A24252 Distribution	12.5 ng	29,197 29,662 29,717	29,525 ± 286	81	19
Availability Codes Dist Avail and/or Special	6.25 ng	32,486 34,744 34,903	34,044 ± 1,352	93	7
AI		1			

Table 1 (concluded)

	3.125 ng	35,150 37,819 37,923	36,964 ± 1,572	100	0
	1.56 ng	36,002 37,475 36,137	36,538 ± 814	100	0
PDGF A (53-63)	500 µg	31,288 31,691 32,193	31,724 ± 453	87	13
	250 µg	36,764 37,412 27,440	33,872 ± 5,580	83	17
	125 µg	37,063 37,758 38,405	37,742 ± 671	92	8
PDGF A (66-78)	500 µg	39,403 37,505 37,546	38,151 ± 1,084	93	7
	250 µg	39,071 41,365 39,426	39,954 ± 1,235	98	2
	125 µg	40,465 40,564	35,349 ±	91	9
PDGF A (77-88)	500 μg	40,071 37,691 38,879	38,880 ± 1,190	100	0
	250 µg	38,983 38,805 39,532	39,107 ± 379	100	0
	125 µg	38,631 38,176 37,584	38,131 ± 523	98	2
PDGF B (25-42)	500 µg	32,668 32,415 33,447	32,843 ± 538	80	20
	250 μ g	29,453 31,978 31,483	30,972 ± 1,338	76	24
	125 µg	35,117 34,305 36,278	35,233 ± 992	86	14

Table 2
THYMIDINE INCORPORATION AT PEPTIDE CONCENTRATION (μg/ml)

Peptide	Control (media)	1000	500	250
PDGF A (53-63)	1,607 ± 305	1,137 ± 73	1,427 ± 197	1,761 ± 118
PDGF A (66-78)	1,544 ± 174	•1,613 ± 121	1,759 ± 208	1,532 ± 276
PDGF A (77-88)	1,474 ± 315	1,507 ± 143	1,630 ± 297	1,637 ± 147
PDGF B (25-42)	1,961 ± 144	2,664 ± 943	1,828 ± 218	1,902 ± 209
PDGF	100 ng 10 ng 1 ng Media	11,681 ± 911 3,935 ± 336 1,897 ± 271 1,527 ± 282		

Table 3
CHEMOTACTIC ACTIVITY OF PEPTIDES

Peptide	Concentration (mg or ng/ml)	OD ₆₅₀ ª	S D	Chemotactic Activity
Media		0.015	0.002	-
PDGF A (53-63)	25 0 μg	0.011	0.002	_
PDGF A (66-78)	250 μς	0.023	0.025	_
PDGF A (77-88)	250 μg	0.053	0.010	+
PDGF B (25-42)	250 μg	0.006	0.006	
PD	GF 20 ng	0.030	0.014	+

^aPeptides were considered positive when the OD was 1-1/2 times more than that of the media.

PLANS FOR NEXT QUARTER

During the next quarter we plan to continue the synthesis of peptides and to conduct cellbinding, mitogenesis, and chemotaxis assays.